

Search for the Decay $\tau^- \rightarrow 4\pi^- 3\pi^+(\pi^0)\nu_\tau$

B. Aubert,¹ R. Barate,¹ D. Boutigny,¹ F. Couderc,¹ Y. Karyotakis,¹ J. P. Lees,¹ V. Poireau,¹ V. Tisserand,¹
A. Zghiche,¹ E. Grauges,² A. Palano,³ M. Pappagallo,³ A. Pompili,³ J. C. Chen,⁴ N. D. Qi,⁴ G. Rong,⁴ P. Wang,⁴
Y. S. Zhu,⁴ G. Eigen,⁵ I. Ofte,⁵ B. Stugu,⁵ G. S. Abrams,⁶ M. Battaglia,⁶ A. W. Borgland,⁶ A. B. Breon,⁶
D. N. Brown,⁶ J. Button-Shafer,⁶ R. N. Cahn,⁶ E. Charles,⁶ C. T. Day,⁶ M. S. Gill,⁶ A. V. Gritsan,⁶ Y. Groyzman,⁶
R. G. Jacobsen,⁶ R. W. Kadel,⁶ J. Kadyk,⁶ L. T. Kerth,⁶ Yu. G. Kolomensky,⁶ G. Kukartsev,⁶ G. Lynch,⁶
L. M. Mir,⁶ P. J. Oddone,⁶ T. J. Orimoto,⁶ M. Pripstein,⁶ N. A. Roe,⁶ M. T. Ronan,⁶ W. A. Wenzel,⁶ M. Barrett,⁷
K. E. Ford,⁷ T. J. Harrison,⁷ A. J. Hart,⁷ C. M. Hawkes,⁷ S. E. Morgan,⁷ A. T. Watson,⁷ M. Fritsch,⁸ K. Goetzen,⁸
T. Held,⁸ H. Koch,⁸ B. Lewandowski,⁸ M. Pelizaeus,⁸ K. Peters,⁸ T. Schroeder,⁸ M. Steinke,⁸ J. T. Boyd,⁹
J. P. Burke,⁹ N. Chevalier,⁹ W. N. Cottingham,⁹ M. P. Kelly,⁹ T. Cuhadar-Donszelmann,¹⁰ C. Hearty,¹⁰
N. S. Knecht,¹⁰ T. S. Mattison,¹⁰ J. A. McKenna,¹⁰ A. Khan,¹¹ P. Kyberd,¹¹ L. Teodorescu,¹¹ A. E. Blinov,¹²
V. E. Blinov,¹² A. D. Bukin,¹² V. P. Druzhinin,¹² V. B. Golubev,¹² E. A. Kravchenko,¹² A. P. Onuchin,¹²
S. I. Serednyakov,¹² Yu. I. Skovpen,¹² E. P. Solodov,¹² A. N. Yushkov,¹² D. Best,¹³ M. Bondioli,¹³ M. Bruinsma,¹³
M. Chao,¹³ I. Eschrich,¹³ D. Kirkby,¹³ A. J. Lankford,¹³ M. Mandelkern,¹³ R. K. Mommsen,¹³ W. Roethel,¹³
D. P. Stoker,¹³ C. Buchanan,¹⁴ B. L. Hartfiel,¹⁴ A. J. R. Weinstein,¹⁴ S. D. Foulkes,¹⁵ J. W. Gary,¹⁵ O. Long,¹⁵
B. C. Shen,¹⁵ K. Wang,¹⁵ L. Zhang,¹⁵ D. del Re,¹⁶ H. K. Hadavand,¹⁶ E. J. Hill,¹⁶ D. B. MacFarlane,¹⁶
H. P. Paar,¹⁶ S. Rahatlou,¹⁶ V. Sharma,¹⁶ J. W. Berryhill,¹⁷ C. Campagnari,¹⁷ A. Cunha,¹⁷ B. Dahmes,¹⁷
T. M. Hong,¹⁷ A. Lu,¹⁷ M. A. Mazur,¹⁷ J. D. Richman,¹⁷ W. Verkerke,¹⁷ T. W. Beck,¹⁸ A. M. Eisner,¹⁸
C. J. Flacco,¹⁸ C. A. Heusch,¹⁸ J. Kroseberg,¹⁸ W. S. Lockman,¹⁸ G. Nesom,¹⁸ T. Schalk,¹⁸ B. A. Schumm,¹⁸
A. Seiden,¹⁸ P. Spradlin,¹⁸ D. C. Williams,¹⁸ M. G. Wilson,¹⁸ J. Albert,¹⁹ E. Chen,¹⁹ G. P. Dubois-Felsmann,¹⁹
A. Dvoretzskii,¹⁹ D. G. Hitlin,¹⁹ I. Narsky,¹⁹ T. Piatenko,¹⁹ F. C. Porter,¹⁹ A. Ryd,¹⁹ A. Samuel,¹⁹ R. Andreassen,²⁰
S. Jayatilke,²⁰ G. Mancinelli,²⁰ B. T. Meadows,²⁰ M. D. Sokoloff,²⁰ F. Blanc,²¹ P. Bloom,²¹ S. Chen,²¹
W. T. Ford,²¹ U. Nauenberg,²¹ A. Olivas,²¹ P. Rankin,²¹ W. O. Ruddick,²¹ J. G. Smith,²¹ K. A. Ulmer,²¹
S. R. Wagner,²¹ J. Zhang,²¹ A. Chen,²² E. A. Eckhart,²² A. Soffer,²² W. H. Toki,²² R. J. Wilson,²² Q. Zeng,²²
E. Feltresi,²³ A. Hauke,²³ B. Spaan,²³ D. Altenburg,²⁴ T. Brandt,²⁴ J. Brose,²⁴ M. Dickopp,²⁴ V. Klose,²⁴
H. M. Lacker,²⁴ R. Nogowski,²⁴ S. Otto,²⁴ A. Petzold,²⁴ G. Schott,²⁴ J. Schubert,²⁴ K. R. Schubert,²⁴
R. Schwierz,²⁴ J. E. Sundermann,²⁴ D. Bernard,²⁵ G. R. Bonneaud,²⁵ P. Grenier,²⁵ S. Schrenk,²⁵ Ch. Thiebaux,²⁵
G. Vasileiadis,²⁵ M. Verderi,²⁵ D. J. Bard,²⁶ P. J. Clark,²⁶ W. Gradl,²⁶ F. Muheim,²⁶ S. Playfer,²⁶ Y. Xie,²⁶
M. Andreotti,²⁷ V. Azzolini,²⁷ D. Bettoni,²⁷ C. Bozzi,²⁷ R. Calabrese,²⁷ G. Cibinetto,²⁷ E. Luppi,²⁷ M. Negrini,²⁷
L. Piemontese,²⁷ F. Anulli,²⁸ R. Baldini-Ferrolì,²⁸ A. Calcaterra,²⁸ R. de Sangro,²⁸ G. Finocchiaro,²⁸ P. Patteri,²⁸
I. M. Peruzzi,²⁸ M. Piccolo,²⁸ A. Zallo,²⁸ A. Buzzo,²⁹ R. Capra,²⁹ R. Contri,²⁹ M. Lo Vetere,²⁹ M. Macri,²⁹
M. R. Monge,²⁹ S. Passaggio,²⁹ C. Patrignani,²⁹ E. Robutti,²⁹ A. Santroni,²⁹ S. Tosi,²⁹ S. Bailey,³⁰
G. Brandenburg,³⁰ K. S. Chaisanguanthum,³⁰ M. Morii,³⁰ E. Won,³⁰ R. S. Dubitzky,³¹ U. Langenegger,³¹
J. Marks,³¹ S. Schenk,³¹ U. Uwer,³¹ W. Bhimji,³² D. A. Bowerman,³² P. D. Dauncey,³² U. Egede,³² R. L. Flack,³²
J. R. Gaillard,³² G. W. Morton,³² J. A. Nash,³² M. B. Nikolich,³² G. P. Taylor,³² M. J. Charles,³³ W. F. Mader,³³
U. Mallik,³³ A. K. Mohapatra,³³ J. Cochran,³⁴ H. B. Crawley,³⁴ V. Eyges,³⁴ W. T. Meyer,³⁴ S. Prell,³⁴
E. I. Rosenberg,³⁴ A. E. Rubin,³⁴ J. Yi,³⁴ N. Arnaud,³⁵ M. Davier,³⁵ X. Giroux,³⁵ G. Grosdidier,³⁵ A. Höcker,³⁵
F. Le Diberder,³⁵ V. Lepeltier,³⁵ A. M. Lutz,³⁵ A. Oyanguren,³⁵ T. C. Petersen,³⁵ M. Pierini,³⁵ S. Plaszczynski,³⁵
S. Rodier,³⁵ P. Roudeau,³⁵ M. H. Schune,³⁵ A. Stocchi,³⁵ G. Wormser,³⁵ C. H. Cheng,³⁶ D. J. Lange,³⁶
M. C. Simani,³⁶ D. M. Wright,³⁶ A. J. Bevan,³⁷ C. A. Chavez,³⁷ J. P. Coleman,³⁷ I. J. Forster,³⁷ J. R. Fry,³⁷
E. Gabathuler,³⁷ R. Gamet,³⁷ K. A. George,³⁷ D. E. Hutchcroft,³⁷ R. J. Parry,³⁷ D. J. Payne,³⁷ K. C. Schofield,³⁷
C. Touramanis,³⁷ C. M. Cormack,³⁸ F. Di Lodovico,³⁸ R. Sacco,³⁸ C. L. Brown,³⁹ G. Cowan,³⁹ H. U. Flaecher,³⁹
M. G. Green,³⁹ D. A. Hopkins,³⁹ P. S. Jackson,³⁹ T. R. McMahon,³⁹ S. Ricciardi,³⁹ F. Salvatore,³⁹ D. Brown,⁴⁰
C. L. Davis,⁴⁰ J. Allison,⁴¹ N. R. Barlow,⁴¹ R. J. Barlow,⁴¹ M. C. Hodgkinson,⁴¹ G. D. Lafferty,⁴¹ M. T. Naisbit,⁴¹
J. C. Williams,⁴¹ C. Chen,⁴² A. Farbin,⁴² W. D. Hulsbergen,⁴² A. Jawahery,⁴² D. Kovalskyi,⁴² C. K. Lae,⁴²
V. Lillard,⁴² D. A. Roberts,⁴² G. Simi,⁴² G. Blaylock,⁴³ C. Dallapiccola,⁴³ S. S. Hertzbach,⁴³ R. Kofler,⁴³
V. B. Koptchev,⁴³ X. Li,⁴³ T. B. Moore,⁴³ S. Saremi,⁴³ H. Staengle,⁴³ S. Willocq,⁴³ R. Cowan,⁴⁴ K. Koeneke,⁴⁴
G. Sciolla,⁴⁴ S. J. Sekula,⁴⁴ F. Taylor,⁴⁴ R. K. Yamamoto,⁴⁴ H. Kim,⁴⁵ P. M. Patel,⁴⁵ S. H. Robertson,⁴⁵
A. Lazzaro,⁴⁶ V. Lombardo,⁴⁶ F. Palombo,⁴⁶ J. M. Bauer,⁴⁷ L. Cremaldi,⁴⁷ V. Eschenburg,⁴⁷ R. Godang,⁴⁷
R. Kroeger,⁴⁷ J. Reidy,⁴⁷ D. A. Sanders,⁴⁷ D. J. Summers,⁴⁷ H. W. Zhao,⁴⁷ S. Brunet,⁴⁸ D. Côté,⁴⁸ P. Taras,⁴⁸
B. Viaud,⁴⁸ H. Nicholson,⁴⁹ N. Cavallo,⁵⁰ G. De Nardo,⁵⁰ C. Gatto,⁵⁰ L. Lista,⁵⁰ D. Monorchio,⁵⁰ P. Paolucci,⁵⁰

D. Piccolo,⁵⁰ C. Sciacca,⁵⁰ M. Baak,⁵¹ H. Bulten,⁵¹ G. Raven,⁵¹ H. L. Snoek,⁵¹ L. Wilden,⁵¹ C. P. Jessop,⁵² J. M. LoSecco,⁵² T. Allmendinger,⁵³ G. Benelli,⁵³ K. K. Gan,⁵³ K. Honscheid,⁵³ D. Hufnagel,⁵³ P. D. Jackson,⁵³ H. Kagan,⁵³ R. Kass,⁵³ T. Pulliam,⁵³ A. M. Rahimi,⁵³ R. Ter-Antonyan,⁵³ Q. K. Wong,⁵³ J. Brau,⁵⁴ R. Frey,⁵⁴ O. Igonkina,⁵⁴ M. Lu,⁵⁴ C. T. Potter,⁵⁴ N. B. Sinev,⁵⁴ D. Strom,⁵⁴ E. Torrence,⁵⁴ F. Colecchia,⁵⁵ A. Dorigo,⁵⁵ F. Galeazzi,⁵⁵ M. Margoni,⁵⁵ M. Morandin,⁵⁵ M. Posocco,⁵⁵ M. Rotondo,⁵⁵ F. Simonetto,⁵⁵ R. Stroili,⁵⁵ C. Voci,⁵⁵ M. Benayoun,⁵⁶ H. Briand,⁵⁶ J. Chauveau,⁵⁶ P. David,⁵⁶ L. Del Buono,⁵⁶ Ch. de la Vaissière,⁵⁶ O. Hamon,⁵⁶ M. J. J. John,⁵⁶ Ph. Leruste,⁵⁶ J. Malclès,⁵⁶ J. Ocariz,⁵⁶ L. Roos,⁵⁶ G. Therin,⁵⁶ P. K. Behera,⁵⁷ L. Gladney,⁵⁷ Q. H. Guo,⁵⁷ J. Panetta,⁵⁷ M. Biasini,⁵⁸ R. Covarelli,⁵⁸ S. Pacetti,⁵⁸ M. Pioppi,⁵⁸ C. Angelini,⁵⁹ G. Batignani,⁵⁹ S. Bettarini,⁵⁹ F. Bucci,⁵⁹ G. Calderini,⁵⁹ M. Carpinelli,⁵⁹ R. Cenci,⁵⁹ F. Forti,⁵⁹ M. A. Giorgi,⁵⁹ A. Lusiani,⁵⁹ G. Marchiori,⁵⁹ M. Morganti,⁵⁹ N. Neri,⁵⁹ E. Paoloni,⁵⁹ M. Rama,⁵⁹ G. Rizzo,⁵⁹ J. Walsh,⁵⁹ M. Haire,⁶⁰ D. Judd,⁶⁰ K. Paick,⁶⁰ D. E. Wagoner,⁶⁰ J. Biesiada,⁶¹ N. Danielson,⁶¹ P. Elmer,⁶¹ Y. P. Lau,⁶¹ C. Lu,⁶¹ J. Olsen,⁶¹ A. J. S. Smith,⁶¹ A. V. Telnov,⁶¹ F. Bellini,⁶² G. Cavoto,⁶² A. D'Orazio,⁶² E. Di Marco,⁶² R. Faccini,⁶² F. Ferrarotto,⁶² F. Ferroni,⁶² M. Gaspero,⁶² L. Li Gioi,⁶² M. A. Mazzoni,⁶² S. Morganti,⁶² G. Piredda,⁶² F. Polci,⁶² F. Safai Tehrani,⁶² C. Voena,⁶² H. Schröder,⁶³ G. Wagner,⁶³ R. Waldi,⁶³ T. Adye,⁶⁴ N. De Groot,⁶⁴ B. Franek,⁶⁴ G. P. Gopal,⁶⁴ E. O. Olaiya,⁶⁴ F. F. Wilson,⁶⁴ R. Aleksan,⁶⁵ S. Emery,⁶⁵ A. Gaidot,⁶⁵ S. F. Ganzhur,⁶⁵ P.-F. Giraud,⁶⁵ G. Graziani,⁶⁵ G. Hamel de Monchenault,⁶⁵ W. Kozanecki,⁶⁵ M. Legendre,⁶⁵ G. W. London,⁶⁵ B. Mayer,⁶⁵ G. Vasseur,⁶⁵ Ch. Yèche,⁶⁵ M. Zito,⁶⁵ M. V. Purohit,⁶⁶ A. W. Weidemann,⁶⁶ J. R. Wilson,⁶⁶ F. X. Yumiceva,⁶⁶ T. Abe,⁶⁷ M. T. Allen,⁶⁷ D. Aston,⁶⁷ R. Bartoldus,⁶⁷ N. Berger,⁶⁷ A. M. Boyarski,⁶⁷ O. L. Buchmueller,⁶⁷ R. Claus,⁶⁷ M. R. Convery,⁶⁷ M. Cristinziani,⁶⁷ J. C. Dingfelder,⁶⁷ D. Dong,⁶⁷ J. Dorfan,⁶⁷ D. Dujmic,⁶⁷ W. Dunwoodie,⁶⁷ S. Fan,⁶⁷ R. C. Field,⁶⁷ T. Glanzman,⁶⁷ S. J. Gowdy,⁶⁷ T. Hadig,⁶⁷ V. Halyo,⁶⁷ C. Hast,⁶⁷ T. Hryn'ova,⁶⁷ W. R. Innes,⁶⁷ M. H. Kelsey,⁶⁷ P. Kim,⁶⁷ M. L. Kocian,⁶⁷ D. W. G. S. Leith,⁶⁷ J. Libby,⁶⁷ S. Luitz,⁶⁷ V. Luth,⁶⁷ H. L. Lynch,⁶⁷ H. Marsiske,⁶⁷ R. Messner,⁶⁷ D. R. Muller,⁶⁷ C. P. O'Grady,⁶⁷ V. E. Ozcan,⁶⁷ A. Perazzo,⁶⁷ M. Perl,⁶⁷ B. N. Ratcliff,⁶⁷ A. Roodman,⁶⁷ A. A. Salnikov,⁶⁷ R. H. Schindler,⁶⁷ J. Schwiening,⁶⁷ A. Snyder,⁶⁷ J. Stelzer,⁶⁷ J. Strube,⁶⁷ D. Su,⁶⁷ M. K. Sullivan,⁶⁷ K. Suzuki,⁶⁷ S. Swain,⁶⁷ J. M. Thompson,⁶⁷ J. Va'vra,⁶⁷ M. Weaver,⁶⁷ W. J. Wisniewski,⁶⁷ M. Wittgen,⁶⁷ D. H. Wright,⁶⁷ A. K. Yarritu,⁶⁷ K. Yi,⁶⁷ C. C. Young,⁶⁷ P. R. Burchat,⁶⁸ A. J. Edwards,⁶⁸ S. A. Majewski,⁶⁸ B. A. Petersen,⁶⁸ C. Roat,⁶⁸ M. Ahmed,⁶⁹ S. Ahmed,⁶⁹ M. S. Alam,⁶⁹ J. A. Ernst,⁶⁹ M. A. Saeed,⁶⁹ M. Saleem,⁶⁹ F. R. Wappler,⁶⁹ S. B. Zain,⁶⁹ W. Bugg,⁷⁰ M. Krishnamurthy,⁷⁰ S. M. Spanier,⁷⁰ R. Eckmann,⁷¹ J. L. Ritchie,⁷¹ A. Satpathy,⁷¹ R. F. Schwitters,⁷¹ J. M. Izen,⁷² I. Kitayama,⁷² X. C. Lou,⁷² S. Ye,⁷² F. Bianchi,⁷³ M. Bona,⁷³ F. Gallo,⁷³ D. Gamba,⁷³ M. Bomben,⁷⁴ L. Bosisio,⁷⁴ C. Cartaro,⁷⁴ F. Cossutti,⁷⁴ G. Della Ricca,⁷⁴ S. Dittongo,⁷⁴ S. Grancagnolo,⁷⁴ L. Lanceri,⁷⁴ P. Poropat,⁷⁴ * L. Vitale,⁷⁴ F. Martinez-Vidal,⁷⁵ R. S. Panvini,⁷⁶ * Sw. Banerjee,⁷⁷ B. Bhuyan,⁷⁷ C. M. Brown,⁷⁷ D. Fortin,⁷⁷ K. Hamano,⁷⁷ R. Kowalewski,⁷⁷ J. M. Roney,⁷⁷ R. J. Sobie,⁷⁷ J. J. Back,⁷⁸ P. F. Harrison,⁷⁸ T. E. Latham,⁷⁸ G. B. Mohanty,⁷⁸ H. R. Band,⁷⁹ X. Chen,⁷⁹ B. Cheng,⁷⁹ S. Dasu,⁷⁹ M. Datta,⁷⁹ A. M. Eichenbaum,⁷⁹ K. T. Flood,⁷⁹ M. Graham,⁷⁹ J. J. Hollar,⁷⁹ J. R. Johnson,⁷⁹ P. E. Kutter,⁷⁹ H. Li,⁷⁹ R. Liu,⁷⁹ B. Mellado,⁷⁹ A. Mihalyi,⁷⁹ Y. Pan,⁷⁹ R. Prepost,⁷⁹ P. Tan,⁷⁹ J. H. von Wimmersperg-Toeller,⁷⁹ J. Wu,⁷⁹ S. L. Wu,⁷⁹ Z. Yu,⁷⁹ M. G. Greene,⁸⁰ H. Neal,⁸⁰ and F. Fabozzi⁸¹

(The BABAR Collaboration)

¹Laboratoire de Physique des Particules, F-74941 Annecy-le-Vieux, France

²IFAE, Universitat Autònoma de Barcelona, E-08193 Bellaterra, Barcelona, Spain

³Università di Bari, Dipartimento di Fisica and INFN, I-70126 Bari, Italy

⁴Institute of High Energy Physics, Beijing 100039, China

⁵University of Bergen, Inst. of Physics, N-5007 Bergen, Norway

⁶Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA

⁷University of Birmingham, Birmingham, B15 2TT, United Kingdom

⁸Ruhr Universität Bochum, Institut für Experimentalphysik 1, D-44780 Bochum, Germany

⁹University of Bristol, Bristol BS8 1TL, United Kingdom

¹⁰University of British Columbia, Vancouver, British Columbia, Canada V6T 1Z1

¹¹Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom

¹²Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia

¹³University of California at Irvine, Irvine, California 92697, USA

¹⁴University of California at Los Angeles, Los Angeles, California 90024, USA

¹⁵University of California at Riverside, Riverside, California 92521, USA

¹⁶University of California at San Diego, La Jolla, California 92093, USA

¹⁷University of California at Santa Barbara, Santa Barbara, California 93106, USA

¹⁸University of California at Santa Cruz, Institute for Particle Physics, Santa Cruz, California 95064, USA

- ¹⁹California Institute of Technology, Pasadena, California 91125, USA
²⁰University of Cincinnati, Cincinnati, Ohio 45221, USA
²¹University of Colorado, Boulder, Colorado 80309, USA
²²Colorado State University, Fort Collins, Colorado 80523, USA
²³Universität Dortmund, Institut für Physik, D-44221 Dortmund, Germany
²⁴Technische Universität Dresden, Institut für Kern- und Teilchenphysik, D-01062 Dresden, Germany
²⁵Ecole Polytechnique, LLR, F-91128 Palaiseau, France
²⁶University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom
²⁷Università di Ferrara, Dipartimento di Fisica and INFN, I-44100 Ferrara, Italy
²⁸Laboratori Nazionali di Frascati dell'INFN, I-00044 Frascati, Italy
²⁹Università di Genova, Dipartimento di Fisica and INFN, I-16146 Genova, Italy
³⁰Harvard University, Cambridge, Massachusetts 02138, USA
³¹Universität Heidelberg, Physikalisches Institut, Philosophenweg 12, D-69120 Heidelberg, Germany
³²Imperial College London, London, SW7 2AZ, United Kingdom
³³University of Iowa, Iowa City, Iowa 52242, USA
³⁴Iowa State University, Ames, Iowa 50011-3160, USA
³⁵Laboratoire de l'Accélérateur Linéaire, F-91898 Orsay, France
³⁶Lawrence Livermore National Laboratory, Livermore, California 94550, USA
³⁷University of Liverpool, Liverpool L69 7ZE, United Kingdom
³⁸Queen Mary, University of London, E1 4NS, United Kingdom
³⁹University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United Kingdom
⁴⁰University of Louisville, Louisville, Kentucky 40292, USA
⁴¹University of Manchester, Manchester M13 9PL, United Kingdom
⁴²University of Maryland, College Park, Maryland 20742, USA
⁴³University of Massachusetts, Amherst, Massachusetts 01003, USA
⁴⁴Massachusetts Institute of Technology, Laboratory for Nuclear Science, Cambridge, Massachusetts 02139, USA
⁴⁵McGill University, Montréal, Quebec, Canada H3A 2T8
⁴⁶Università di Milano, Dipartimento di Fisica and INFN, I-20133 Milano, Italy
⁴⁷University of Mississippi, University, Mississippi 38677, USA
⁴⁸Université de Montréal, Laboratoire René J. A. Lévêque, Montréal, Quebec, Canada H3C 3J7
⁴⁹Mount Holyoke College, South Hadley, Massachusetts 01075, USA
⁵⁰Università di Napoli Federico II, Dipartimento di Scienze Fisiche and INFN, I-80126, Napoli, Italy
⁵¹NIKHEF, National Institute for Nuclear Physics and High Energy Physics, NL-1009 DB Amsterdam, The Netherlands
⁵²University of Notre Dame, Notre Dame, Indiana 46556, USA
⁵³Ohio State University, Columbus, Ohio 43210, USA
⁵⁴University of Oregon, Eugene, Oregon 97403, USA
⁵⁵Università di Padova, Dipartimento di Fisica and INFN, I-35131 Padova, Italy
⁵⁶Universités Paris VI et VII, Laboratoire de Physique Nucléaire et de Hautes Energies, F-75252 Paris, France
⁵⁷University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA
⁵⁸Università di Perugia, Dipartimento di Fisica and INFN, I-06100 Perugia, Italy
⁵⁹Università di Pisa, Dipartimento di Fisica, Scuola Normale Superiore and INFN, I-56127 Pisa, Italy
⁶⁰Prairie View A&M University, Prairie View, Texas 77446, USA
⁶¹Princeton University, Princeton, New Jersey 08544, USA
⁶²Università di Roma La Sapienza, Dipartimento di Fisica and INFN, I-00185 Roma, Italy
⁶³Universität Rostock, D-18051 Rostock, Germany
⁶⁴Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX, United Kingdom
⁶⁵DSM/Dapnia, CEA/Saclay, F-91191 Gif-sur-Yvette, France
⁶⁶University of South Carolina, Columbia, South Carolina 29208, USA
⁶⁷Stanford Linear Accelerator Center, Stanford, California 94309, USA
⁶⁸Stanford University, Stanford, California 94305-4060, USA
⁶⁹State University of New York, Albany, New York 12222, USA
⁷⁰University of Tennessee, Knoxville, Tennessee 37996, USA
⁷¹University of Texas at Austin, Austin, Texas 78712, USA
⁷²University of Texas at Dallas, Richardson, Texas 75083, USA
⁷³Università di Torino, Dipartimento di Fisica Sperimentale and INFN, I-10125 Torino, Italy
⁷⁴Università di Trieste, Dipartimento di Fisica and INFN, I-34127 Trieste, Italy
⁷⁵IFIC, Universitat de Valencia-CSIC, E-46071 Valencia, Spain
⁷⁶Vanderbilt University, Nashville, Tennessee 37235, USA
⁷⁷University of Victoria, Victoria, British Columbia, Canada V8W 3P6
⁷⁸Department of Physics, University of Warwick, Coventry CV4 7AL, United Kingdom
⁷⁹University of Wisconsin, Madison, Wisconsin 53706, USA
⁸⁰Yale University, New Haven, Connecticut 06511, USA
⁸¹Università della Basilicata, I-85100 Potenza, Italy

(Dated: February 7, 2008)

A search for the decay of the τ lepton to seven charged pions and one or zero π^0 mesons was performed using the *BABAR* detector at the PEP-II asymmetric-energy e^+e^- collider. The analysis uses 232.2 fb^{-1} of data at center-of-mass energies on or near the $\Upsilon(4S)$ resonance. We observe 24 events with an expected background of 21.6 ± 1.3 events. Without evidence for a signal, we calculate an upper limit of $\mathcal{B}(\tau^- \rightarrow 4\pi^- 3\pi^+(\pi^0)\nu_\tau) < 3.0 \times 10^{-7}$ at 90 % confidence level. This is an improvement by nearly an order of magnitude over the previously established limit. In addition, we set upper limits for the exclusive decays $\tau^- \rightarrow 4\pi^- 3\pi^+\nu_\tau$ and $\tau^- \rightarrow 4\pi^- 3\pi^+\pi^0\nu_\tau$.

PACS numbers: 13.35.Dx, 13.66.De, 13.85.Rm

The decay of the τ lepton to seven charged particles [12] has not been observed to date. An upper limit of $\mathcal{B}(\tau^- \rightarrow 4\pi^- 3\pi^+(\pi^0)\nu_\tau) < 2.4 \times 10^{-6}$ at 90 % confidence level has been set by the CLEO Collaboration [1]. Theoretical calculations using an effective chiral Lagrangian to estimate the matrix elements show that the τ decay rate to seven charged particles is much smaller than the decay into five charged particles due to its smaller phase space. This leads to a theoretical branching fraction estimate of the order of 10^{-11} , which could be enhanced by up to an order of magnitude if this decay proceeds via resonances [2].

This analysis is based on data recorded by the *BABAR* detector at the PEP-II asymmetric-energy e^+e^- storage ring operated at the Stanford Linear Accelerator Center. The data sample consists of 232.2 fb^{-1} recorded at center-of-mass (CM) energies of 10.58 GeV and 10.54 GeV. With an expected cross section for τ pairs for these CM energies of $\sigma_{\tau\tau} = (0.89 \pm 0.02)\text{ nb}$ [3], the number of produced τ pair events $N_{\tau\tau}$ is $(206.6 \pm 5.2) \times 10^6$.

The *BABAR* detector is described in detail in Ref. [4] and only a brief description is given here. Charged-particle momenta are measured with a 5-layer double-sided silicon vertex tracker (SVT) and a 40-layer drift chamber (DCH) inside a 1.5-T solenoidal magnetic field. A calorimeter (EMC) consisting of 6580 CsI(Tl) crystals is used to measure electromagnetic energy. A ring-imaging Cherenkov detector is used to identify charged hadrons, in combination with ionization energy loss measurements in the SVT and the DCH. Muons are identified by an instrumented magnetic-flux return (IFR). Particle four-momenta are reconstructed in the laboratory frame and then boosted to the e^+e^- center-of-mass (CM) frame using the measured beam energies.

We use Monte Carlo simulation techniques to estimate our signal efficiencies and background contamination from other τ -pair events. The production of τ pairs is simulated with the KK generator [5], and non-signal τ lepton decays are modeled with TAUOLA [6] according to measured rates [7]. Signal events were generated with uniform density throughout the available phase space. The simulation of the *BABAR* detector is based on GEANT 4 [8].

Events with eight charged tracks and a net charge of zero are selected. Tracks are required to have a distance of closest approach to the interaction point in the plane transverse to the beam axis (DOCA_{XY}) of less than 1.5 cm, and a distance of closest approach along the beam direction of less than 10 cm. It is required that at least five of these eight tracks have a minimum transverse momentum (p_T) of 100 MeV/ c , and 12 or more DCH hits. Photons are reconstructed from EMC clusters and are required to have a minimum energy of 70 MeV, more than three crystal hits, and a lateral energy profile consistent with that of a photon. The event is divided into hemispheres by a plane perpendicular to the thrust axis [9], where the thrust is calculated from all charged tracks and all photons in the event. The event thrust has to be larger than 0.9. We require all events to have one track in one hemisphere (the “tag-side” hemisphere) and seven tracks in the other hemisphere (the “signal side” hemisphere). The above requirements define the 1-7 topology.

Monte Carlo studies have shown that the background from τ -pair events stems from photon conversions in the detector material. To reduce this contribution and to exclude $e^+e^- \rightarrow q\bar{q}$ ($q = \{u, d, s, c, b\}$) decays containing kaons we apply particle identification on the signal side. We demand that at least six of the tracks are identified as pions with high probability. We apply looser identification criteria to the seventh track.

To further reduce photon conversions, we require all seven tracks on the signal side to have a minimum transverse momentum of 100 MeV/ c and the ratio of DOCA_{XY}/p_T to be less than $0.7\text{ cm}/(\text{GeV}/c)$.

Further suppression of $e^+e^- \rightarrow q\bar{q}$ events is achieved by requiring the tag side to satisfy one of the following criteria: 1) a tightly identified electron or muon with no more than one additional photon on the tag-side; 2) a tight lepton veto and no additional photon; or 3) a reconstructed ρ meson with an invariant mass of $0.650 < m_\rho < 0.875\text{ GeV}/c^2$, derived from the combination of the 1-prong track with a reconstructed π^0 candidate. The π^0 candidates are formed by combining two photons and requiring an invariant mass between 0.113 and 0.155 GeV/c^2 .

After applying these requirements we calculate the

TABLE I: Cumulative signal efficiencies in % and τ background contributions as number of events scaled to the data luminosity after the various selection criteria. ID denotes Identification, Conv. Conversion, and Gen. Generic.

	Efficiency [%]		Number of events		
	$7\pi\nu_\tau$	$7\pi\pi^0\nu_\tau$	Gen. τ	$5\pi\nu_\tau$	$5\pi\pi^0\nu_\tau$
1-7 Topology	23.6 ± 1.4	22.1 ± 1.4	767	198	187
Particle ID	20.7 ± 1.3	19.6 ± 1.3	108	64	75
Conv. Veto	15.8 ± 1.0	14.9 ± 1.0	0	4.7	9.2
1-Prong Tag	10.2 ± 0.7	9.6 ± 0.7	0	1.7	4.2
Signal Region	9.4 ± 0.6	9.3 ± 0.6	0	0.4	0.8

pseudo mass (m) of the τ lepton [10]:

$$m^2 = 2(E_{beam} - E_{7\pi})(E_{7\pi} - P_{7\pi}) + m_{7\pi}^2. \quad (1)$$

The pseudo mass is an approximation of the invariant mass of the tau, where the neutrino's flight direction is approximated by the combined momentum vector $P_{7\pi}$ of the seven charged tracks, and its energy is taken to be the difference between the beam energy E_{beam} in the center-of-mass system and the combined energy $E_{7\pi}$ of the charged tracks. The pseudo mass allows for a better discrimination between signal events and background from $e^+e^- \rightarrow q\bar{q}$ events than $m_{7\pi}$. The final event count is performed in the signal region $1.3 < m < 1.8 \text{ GeV}/c^2$. Figure 1 illustrates the pseudo mass spectra of simulated signal and background contributions after the topology selection.

For this analysis, a comparison of Monte Carlo simulation and data has shown that $e^+e^- \rightarrow q\bar{q}$ background contributions cannot reliably be extracted from simulation due to difficulties in modeling the fragmentation processes. The shape of the pseudo mass distribution appears to be correctly modeled, but the overall normalization is not. Therefore, we estimate this background directly from data, using a method described below. On the other hand, the simulation of τ pair events yields a reliable estimate of their expected background contribution, verified by loosening requirements that suppress the tau background. A breakdown of the signal efficiencies and individual τ background contributions are listed in Table I for each selection step. In this table the generic τ sample does not contain 5π and $5\pi\pi^0$ decays, which are listed separately since they comprise the only background from τ decays after the final selection.

The efficiencies for events in the signal region for $\tau^- \rightarrow 4\pi^-3\pi^+\nu_\tau$ and $\tau^- \rightarrow 4\pi^-3\pi^+\pi^0\nu_\tau$ are $(9.4\pm0.1\pm0.6)\%$ and $(9.3\pm0.1\pm0.6)\%$ (see Table I). The first errors are statistical and the second systematic.

The systematic errors on the signal efficiencies include contributions from uncertainties in the reconstruction of charged tracks (5.2%), the uncertainty associated with the particle identification on the signal and




FIG. 1: Pseudo mass distributions of the seven charged tracks after the 1-7 topology selection for signal (dashed-dotted: $\tau^- \rightarrow 4\pi^-3\pi^+\pi^0\nu_\tau$; dotted: $\tau^- \rightarrow 4\pi^-3\pi^+\nu_\tau$), generic τ (including 5-prong decays) (dashed), and $e^+e^- \rightarrow q\bar{q}$ Monte Carlo events (solid). Entries are normalized to the data luminosity. To display the two signal modes we assumed a branching fraction of 2×10^{-5} .

tag side (2.7%), the luminosity measurement and the τ pair cross-section determination (2.3%), and the uncertainty on the generic 1-prong τ decay branching ratios (0.5%). Since the efficiencies of the two decay channels to the 7-prong τ decays are in good agreement, we average them and obtain an overall efficiency for the decay $\tau^- \rightarrow 4\pi^-3\pi^+(\pi^0)\nu_\tau$ of $(9.4\pm0.6)\%$ which includes the statistical and systematic contributions described above.

To determine the number of $e^+e^- \rightarrow q\bar{q}$ events in the signal region, we use the following procedure: we histogram the pseudo mass distribution of all data events that satisfy the topology requirement and are in the pseudo mass sideband $1.8 < m < 2.6 \text{ GeV}/c^2$. The contribution of the τ background, determined from the simulation, is subtracted from this histogram, and the resulting distribution is fit with a Gaussian function. We then fix the mean and the width of this Gaussian, and use it to fit the τ -background-subtracted distributions of sideband events for all subsequent cuts, floating only the normalizations. Integrating the area of the resulting Gaussian functions in the signal region $1.3 < m < 1.8 \text{ GeV}/c^2$ yields an estimate of the $e^+e^- \rightarrow q\bar{q}$ background contributions.

Table II lists the number of expected τ and $e^+e^- \rightarrow q\bar{q}$ background events in the signal region after the four selection steps. In Figure 2 (a) – (d) we show the tau-background-subtracted pseudo mass data distribution after the four selection steps. Although entries below $1.8 \text{ GeV}/c^2$ are shown here, these events were hidden during the development of our analysis to avoid experimenter bias. Overlaid in the individual figures are the fit re-

TABLE II: Predicted and observed number of events in the signal region of $1.3 < m < 1.8 \text{ GeV}/c^2$. The τ background yield is obtained from the simulation, and its error reflects finite Monte Carlo statistics. The $e^+e^- \rightarrow q\bar{q}$ yield is determined by fitting the data in the pseudo mass sideband, and its error results from data statistics.

	τ Bg.	$e^+e^- \rightarrow q\bar{q}$ Bg.	Observed
1-7 Topology	128 ± 13	574 ± 21	695
Particle ID	28 ± 6	241 ± 10	244
Conv. Veto	2.4 ± 1.3	119 ± 5	104
1-Prong Tag	1.3 ± 1.0	20.3 ± 0.7	24

TABLE III: Observed and predicted number of $e^+e^- \rightarrow q\bar{q}$ background events in the 1-8 and 1-7 topology data for the different selection criteria.

	1-8 (fixed)	1-8 (free)	1-8 obs.	1-7 (free)
Topology	19.0 ± 2.7	19.0 ± 2.7	23	574 ± 21
Particle Id	12.2 ± 1.6	11.2 ± 2.0	10	222 ± 19
Conv. Veto	2.7 ± 0.3	2.6 ± 1.4	1	126 ± 18
1-Pr. Tag	0.5 ± 0.1	0.3 ± 0.5	0	20.2 ± 7.7

sults to the pseudo mass spectra in the range of 1.8 to 2.6 GeV/c^2 .

After the final selection we observe 24 events in the data (see Figure 3 (a)) with a total number of predicted background from τ and $e^+e^- \rightarrow q\bar{q}$ events of 21.6 ± 1.2 . The statistical error of the $e^+e^- \rightarrow q\bar{q}$ background estimate is derived from the statistical uncertainties of the parameters in the Gaussian fit.

To validate the $e^+e^- \rightarrow q\bar{q}$ background estimation we use 1-8 topology data [13], which has negligible expected signal contributions. We fit the pseudo mass distribution between 1.8 and 2.6 GeV/c^2 , integrate the fit function in the pseudo mass region 1.3 to 1.8 GeV/c^2 , and compare this with the number of events found in data (see Table III, columns two and four). In addition, we repeat the fit with mean and sigma floating (“free” fit) in each individual distribution to compare the expected numbers of events (Table III, column three). As a final cross check, fits with the Gaussian mean and width floating were performed for the 1-7 topology data, and the results are included in Table III, column five. We note the very good agreement of the fits with fixed or floating Gaussian parameters.

The systematic error on the number of expected background events is ± 0.4 events, derived from variations of the fit range and shape of the extrapolation function. The total number of expected background events is 21.6 ± 1.3 .

We calculate the branching fraction of the $\tau^- \rightarrow 4\pi^- 3\pi^+(\pi^0)\nu_\tau$ decay based on the following likelihood

function, which convolves a Poisson distribution with two Gaussian resolution functions, accounting for the uncertainties in the background and in the efficiency:

$$\mathcal{L}(\mathcal{B}; n, \hat{b}, \hat{f}, b, f) = \frac{\mu^n e^{-\mu}}{n!} \frac{1}{2\pi\sigma_b\sigma_f} e^{-\frac{1}{2}\left(\frac{\hat{b}-b}{\sigma_b}\right)^2 - \frac{1}{2}\left(\frac{\hat{f}-f}{\sigma_f}\right)^2}, \quad (2)$$

where \mathcal{B} denotes the branching fraction of $\tau^- \rightarrow 4\pi^- 3\pi^+(\pi^0)\nu_\tau$, $f = 2N_{\tau\tau}\epsilon$, ϵ is the signal efficiency, $\mu = \langle n \rangle = \hat{f}\mathcal{B} + \hat{b}$, with n the number of observed events, and b and σ_b are the expected background and its error. σ_f incorporates the errors on the signal efficiency and the number of τ pair events.

The likelihood function is maximized with respect to the branching fraction \mathcal{B} , \hat{f} and \hat{b} , and the following numerical value for the branching fraction is obtained by MINUIT[11]:

$$\mathcal{B}(\tau^- \rightarrow 4\pi^- 3\pi^+(\pi^0)\nu_\tau) = (0.7^{+1.4}_{-1.3}) \times 10^{-7}.$$

Since we have no evidence for a signal we compute a Bayesian upper limit using a uniform prior in the branching fraction, the background, and the efficiency. This is done by integrating out \hat{f} and \hat{b} in the likelihood function and plotting \mathcal{L} as function of \mathcal{B} . In this way we normalize the distribution to unity and get as the result of this analysis an upper limit at the point where the integral reaches 0.9:

$$\mathcal{B}(\tau^- \rightarrow 4\pi^- 3\pi^+(\pi^0)\nu_\tau) < 3.0 \times 10^{-7} \text{ (at 90 \% CL)}.$$

With the same approach, setting the number of observed events N_{obs} to the expected number of background events of $N_{exp} = 21.6$, we calculate the sensitivity of the analysis to be $\mathcal{B}^{N_{obs} \equiv N_{exp}}(\tau^- \rightarrow 4\pi^- 3\pi^+(\pi^0)\nu_\tau) < 2.5 \times 10^{-7}$ at 90 % CL.

In addition to this inclusive result, we set limits on the branching fractions of the exclusive decay modes $\tau^- \rightarrow 4\pi^- 3\pi^+\nu_\tau$ and $\tau^- \rightarrow 4\pi^- 3\pi^+\pi^0\nu_\tau$. To select $\tau^- \rightarrow 4\pi^- 3\pi^+\nu_\tau$ candidates in the inclusive sample, we require the number of photons on the signal side to be zero. This yields a signal efficiency for $\tau^- \rightarrow 4\pi^- 3\pi^+\nu_\tau$ decays of $(5.5 \pm 0.3)\%$ while reducing the expected generic τ decay and $e^+e^- \rightarrow q\bar{q}$ background to 3.9 ± 0.8 events. The decays $\tau^- \rightarrow 4\pi^- 3\pi^+\pi^0\nu_\tau$ are treated as background in this case and have a reconstruction efficiency of $(0.8 \pm 0.1)\%$. Reversing this selection by demanding at least one reconstructed π^0 on the signal side yields a signal efficiency for the $\tau^- \rightarrow 4\pi^- 3\pi^+\pi^0\nu_\tau$ decay of $(3.6 \pm 0.3)\%$ and an expected generic τ decay and $e^+e^- \rightarrow q\bar{q}$ background of 8.2 ± 0.5 events. In this case, the reconstruction efficiency for $\tau^- \rightarrow 4\pi^- 3\pi^+\nu_\tau$ is $(0.3 \pm 0.0)\%$. The systematic uncertainties of the $\tau^- \rightarrow 4\pi^- 3\pi^+\nu_\tau$ mode are identical to the inclusive measurement already discussed above. For the




FIG. 2: Pseudo mass distributions for 1-7 topology data. (a) after 1-7 topology selection; (b) after particle identification; (c) after conversion veto; (d) after 1-prong tag.

$\tau^- \rightarrow 4\pi^- 3\pi^+ \pi^0 \nu_\tau$ mode, an additional uncertainty of 5.0 % on the efficiency of the π^0 reconstruction is taken into account.

The likelihood function (2) was modified to accommodate one exclusive mode acting as background for the other. We observe eight events in the $\tau^- \rightarrow 4\pi^- 3\pi^+ \nu_\tau$ signal region and seven events in $\tau^- \rightarrow 4\pi^- 3\pi^+ \pi^0 \nu_\tau$. We calculate the following numerical values for the branching fractions:

$$\begin{aligned} \mathcal{B}(\tau^- \rightarrow 4\pi^- 3\pi^+ \nu_\tau) &= (2.0^{+1.5}_{-1.2}) \times 10^{-7} \\ \mathcal{B}(\tau^- \rightarrow 4\pi^- 3\pi^+ \pi^0 \nu_\tau) &= (-1.0 \pm 1.8) \times 10^{-7}. \end{aligned}$$

Without evidence for a signal we compute Bayesian upper limits using uniform priors in the branching fractions, the backgrounds, and the efficiencies, of

$$\begin{aligned} \mathcal{B}(\tau^- \rightarrow 4\pi^- 3\pi^+ \nu_\tau) &< 4.3 \times 10^{-7} \text{ (at 90 \% CL)} \\ \mathcal{B}(\tau^- \rightarrow 4\pi^- 3\pi^+ \pi^0 \nu_\tau) &< 2.5 \times 10^{-7} \text{ (at 90 \% CL)}. \end{aligned}$$

With the same approach, setting the number of observed events N_{obs} to the expected number of background events, we calculate the sensitivities $\mathcal{B}^{N_{obs} \equiv N_{exp}}(\tau^- \rightarrow 4\pi^- 3\pi^+ \nu_\tau) < 2.2 \times 10^{-7}$ and $\mathcal{B}^{N_{obs} \equiv N_{exp}}(\tau^- \rightarrow 4\pi^- 3\pi^+ \pi^0 \nu_\tau) < 4.2 \times 10^{-7}$. Figures 3 (b) and (c) show details of the data pseudo mass spectra with an overlay of the expected background distributions.

This analysis improves the existing experimental limits by an order of magnitude for the inclusive mode, but is still several orders of magnitude larger than the theoretical prediction. The exclusive decays are reported for the first time and are consistent with the inclusive result.

We are grateful for the extraordinary contributions of our PEP-II colleagues in achieving the excellent luminosity and machine conditions that have made this work possible. The success of this project also relies critically on the expertise and dedication of the computing organizations that support *BABAR*. The collaborating institutions wish to thank SLAC for its support and the kind hospitality extended to them. This work is supported by the US Department of Energy and National Science Foundation, the Natural Sciences and Engineering Research Council (Canada), Institute of High Energy Physics (China), the Commissariat à l'Energie Atomique and Institut National de Physique Nucléaire et de Physique des Particules (France), the Bundesministerium für Bildung und Forschung and Deutsche Forschungsgemeinschaft (Germany), the Istituto Nazionale di Fisica Nucleare (Italy), the Foundation for Fundamental Research on Matter (The Netherlands), the Research Council of Norway, the Ministry of Science and Technology of the Russian Federation, and the Particle Physics and Astronomy Research Council (United Kingdom). Individu-




FIG. 3: Details of the data pseudo mass spectra. (a) $\tau^- \rightarrow 4\pi^-3\pi^+(\pi^0)\nu_\tau$, (b) $\tau^- \rightarrow 4\pi^-3\pi^+\nu_\tau$, (c) $\tau^- \rightarrow 4\pi^-3\pi^+\pi^0\nu_\tau$ event candidates.

als have received support from CONACyT (Mexico), the A. P. Sloan Foundation, the Research Corporation, and the Alexander von Humboldt Foundation.

* Deceased

- [1] K. Edwards *et al.*, Phys. Rev. D **56**, 5297 (1997).
- [2] S. Nussinov and M. V. Purohit, Phys. Rev. **D65**, 034018 (2002).
- [3] B. F. L. Ward, S. Jadach, and Z. Was, Nucl. Phys. Proc. Suppl. **116**, 73 (2003).
- [4] B. Aubert *et al.*, Nucl. Instrum. Meth. **A479**, 1 (2002).
- [5] S. Jadach, B. F. L. Ward, and Z. Was, Comput. Phys. Commun. **130**, 260 (2000).
- [6] S. Jadach, J. H. Kuhn, and Z. Was, Comput. Phys. Commun. **64**, 275 (1990).
- [7] K. Hagiwara *et al.*, Phys. Rev. **D66**, 010001 (2002).
- [8] S. Agostinelli *et al.*, Nucl. Instrum. Meth. **A506**, 250 (2003).
- [9] T. Sjostrand, Comput. Phys. Commun. **39**, 347 (1986).
- [10] H. Albrecht *et al.*, Phys. Lett. B **292**, 221 (1992).
- [11] F. James and M. Roos, Comput. Phys. Commun. **10**, 343 (1975).
- [12] Throughout this paper, whenever a mode is given its charge conjugate is also implied.
- [13] The event selection of the 1-7 topology was modified accordingly, to accommodate eight tracks on the signal side.